Volume 14, 01 November 2014 Publisher: Igitur publishing URL: http://www.ijic.org

Cite this as: Int J Integr Care 2014; Inter Digital Health Suppl; URN:NBN:NL:UI:10-1-116512

Copyright: (cc) BY

Conference Abstract

Self-management in long term conditions using smartphones: A pilot study in Parkinson's disease

Rashmi Naravana. uMotif Digital Health. United Kingdom

Bruce Hellman, uMotif Digital Health, United Kingdom

Caspar Addyman, Birkbeck University of London, United Kingdom

Jon Stamford, The Cure Parkinson's Trust, United Kingdom

Correspondence to: **Rashmi Narayana**, uMotif Digital Health, United Kingdom, E-mail: rashmi@umotif.com

Abstract

Background: Long-term conditions (LTC) or chronic diseases are the leading causes of mortality and morbidity worldwide. Self-management is now an accepted method to manage LTC. Parkinson's disease (PD) is a long-term neurological condition with self-management being a core component of managing the condition optimally.

Aims: The primary aim was to evaluate use of 2 versions of a PD tracker app and assess the impact of use on a range of outcomes including: self-reported measures of adherence to treatment, quality of life, happiness and non-motor symptoms. Qualitative interviews were carried out with a sub-group of participants to assess usability. The study was funded by the Department of Health's Small Business Research Initiative (SBRI).

Methods: 36 patients with PD took part in a 55 day pilot study and were randomised into 2 groups – the limited app group (n=19) received an app which had tools for daily self-tracking on 10 measures of symptom severity, general well-being and health behaviours along with a daily diary; and the full app group (n=17) received an app which had the same tools as the limited app group with the addition of medication reminders and 2 games to assess cognition.

Results: Participants used the app for 55 days, and entered data on at least 70% of those days. There were no statistically significant differences between the two versions of the app on self-reported measures of adherence to treatment, quality of life, happiness and non-motor symptoms. However, there were increases in absolute scores in self-reported measures of adherence to treatment and quality of life. Participants provided positive feedback on the ease of use of the app and value of symptom tracking.

Conclusion: The PD tracker app was used regularly as a self-management tool by patients and could help in improving self-reported measures of adherence and well-being. A larger sample size and stronger study design are needed to confirm these findings conclusively.

Keywords

long-term conditions; self-management; mobile health; parkinson's disease; mobile application

PowerPoint presentation:

https://www.conftool.pro/digital-health-care-2014/index.php?page=adminPapersDetails&path=adminPapers&form_id=80