



Optimising Integrated Stroke Care in Regional Networks: A Nationwide Self-Assessment Study in 2012, 2015 and 2019

RESEARCH AND THEORY

HELENE R. VOOGDT-PRUIS 

NICK ZONNEVELD 

MONIQUE BERGSMA

ELIZE VAN WIJK

HENK KERKHOFF

LUIKJE VAN DER DUSSEN

MAARTJE KUIJPENS

HUBERTUS J. M. VRIJHOEF 

MIRELLA M. N. MINKMAN 

*Author affiliations can be found in the back matter of this article



ABSTRACT

Background: To help enhance the quality of integrated stroke care delivery, regional stroke services networks in the Netherlands participated in a self-assessment study in 2012, 2015 and 2019.

Methods: Coordinators of the regional stroke services networks filled out an online self-assessment questionnaire in 2012, 2015 and 2019. The questionnaire, which was based on the Development Model for Integrated Care, consisted of 97 questions in nine clusters (themes). Cluster scores were calculated as proportions of the activities implemented. Associations between clusters and features of stroke services were assessed by regression analysis.

Results: The response rate varied from 93.1% (2012) to 85.5% (2019). Over the years, the regional stroke services networks increased in 'size': the median number of organisations involved and the volume of patients per network increased (7 and 499 in 2019, compared to 5 and 364 in 2012). At the same time, fewer coordinators were appointed for more than 1 day a week in 2019 (35.1%) compared to 2012 (45.9%). Between 2012 and 2019, there were statistically significantly more elements implemented in four out of nine clusters: 'Transparent entrepreneurship' (MD = 18.0% F(1) = 10.693, p = 0.001), 'Roles and tasks' (MD = 14.0% F(1) = 9.255, p = 0.003), 'Patient-centeredness' (MD = 12.9% F(1) = 9.255, p = 0.003), and 'Commitment' (MD = 11.2%, F(1) = 4.982, p = 0.028). A statistically significant positive correlation was found for all clusters between implementation of activities and age of the network. In addition, the number of involved organisations is associated with better execution of implemented activities for 'Transparent entrepreneurship', 'Result-focused learning' and 'Quality of care'. Conversely, there are small but negative associations between the volume of patients and implementation rates for 'Interprofessional teamwork' and 'Patient-centredness'.

Conclusion: This long-term analyses of stroke service development in the Netherlands, showed that between 2012 and 2019, integrated care activities within the regional stroke networks increased. Experience in collaboration between organisations within a network benefits the uptake of integrated care activities.

CORRESPONDING AUTHOR:

Helene R. Voogdt-Pruis

University Medical Centre
Utrecht, Julius Global Health,
PO Box 85500, 3508 GA
Utrecht, Netherlands

h.r.pruis@umcutrecht.nl

KEYWORDS:

stroke services; integrated care; care coordination; collaboration; quality improvement; interprofessional teamwork; patient-centredness

TO CITE THIS ARTICLE:

Voogdt-Pruis HR, Zonneveld N, Bergsma M, van Wijk E, Kerkhoff H, van der Dussen L, Kuijpers M, Vrijhoef HJM, Minkman MMN. Optimising Integrated Stroke Care in Regional Networks: A Nationwide Self-Assessment Study in 2012, 2015 and 2019. *International Journal of Integrated Care*, 2021; 21(3): 12, 1–11. DOI: <https://doi.org/10.5334/ijic.5611>

BACKGROUND

Worldwide, stroke is the second leading cause of death and source of 'disability adjusted life years'. The prevalence of stroke is expected to increase due to a growing and ageing population and due to lower stroke case fatality rates associated with better acute ischaemic stroke care and improved recurrent stroke prevention strategies addressing metabolic and behavioural factors [1]. Ischaemic stroke occurs when the blood supply to part of the brain is interrupted or reduced, preventing brain tissue from getting oxygen and nutrients. Less frequently, cerebral haemorrhages occur with similar, often devastating results. Early action and effective treatments can reduce brain damage, complications, and disabilities. Recovery after stroke mainly occurs within the first six months. Stroke patients can experience long-term difficulties in terms of quality of life, social reintegration, life satisfaction and emotional functioning, including depression and anxiety [2]. After a stroke, about 70 percent of patients are discharged back home from the hospital, about 20 percent are referred to rehabilitation centres or nursing homes and about 10 percent die within 30 days [3].

In order to deliver patient-centred integrated stroke care, cooperation and collaboration between professionals, patients and caregivers is of utmost importance. A seamless integration of services across the healthcare and social care interface is required for effective treatment and management of strokes [3]. In addition, integrated stroke care is needed to align healthcare services, to decrease repeated assessments and incomplete or conflicting information about the patient's health status and to reduce duplications in supervision and multiple transaction costs [4]. However, various barriers could hinder the formation and development of such collaboration, for example related to existing administration and regulation, funding, organisational and interorganisational domains, service delivery and work routines [5]. From the patients' perspective, quality of stroke care could be improved by better collaboration between professionals, by providing tailor-made information and by shared decision making [6, 7]. In addition, better preparation of transitions from hospital to home care and tailored support at home is needed [3].

Integration of hospital and primary care services has been a priority of health policy in The Netherlands since the 1990s. Several initiatives focused on the implementation or the investigation of so called 'transmural or integrated care'. Transmural care was defined as care attuned to the needs of the patient, provided on the basis of cooperation and coordination between general and specialised care professionals with shared overall responsibility and the specification of delegated responsibilities [8]. For stroke, first studies on the effectiveness of 'transmural care' in the Netherlands

appeared early 2000. Reported benefits of 'transmural stroke care' six months after stroke were: higher patient satisfaction, higher portion of patients back home, and less volume of home care [9]. Further improvement of quality of 'transmural or integrated stroke care' was the aim of a national improvement project that started in 2002. Based on the Breakthrough Series [10] teams from 23 regions aimed to make 'breakthrough' improvements [11] on specific topics in stroke care. Significantly better results in health outcomes and interprofessional collaboration were attained [12]. The 'sense of urgency' for continuous improvement of integrated stroke care was set. Following this, more registered integrated stroke services emerged. Regional networks of service providers, such as hospitals, geriatric and medical rehabilitation centres, skilled nursing facilities, and primary care providers started to work together in an organised way [13]. These networks were led by a steering board with representatives —professionals and managers— of involved organisations. Needed were professionals and managers with the ability to create collaboration and cooperation across professions and organisations, that are comfortable with distributing responsibilities, and that thoroughly understand the aim of integrated stroke services [14]. Often one professional within the stroke service was appointed as coordinator. The aim was to work together to provide multidisciplinary, coordinated care through organised patient transfers and multidisciplinary protocols. In addition, the ambition was to make continuous improvements on collaboration and cooperation of stroke care. Integrated stroke services in the Netherlands turned out to increase patient satisfaction and the portion of patients back home, as well as reducing the volume of home care [6, 11–14].

In 2006, the Dutch Knowledge network of stroke services (KNCN) was founded to support all stroke services in the Netherlands in their mission to improve the coordination, cooperation and quality of regional multidisciplinary integrated stroke care. In 2006, 21 stroke services participated; this increased to 44 in 2009. In 2012 and 2019, 72 and 52 stroke services were member of KNCN. A decline in participating stroke services took place after 2014 due to the changed role of KNCN in the national Benchmark stroke care and discontinuation of financial support, and due to the merging of some stroke services. Nonetheless, in 2019, about 90% of all hospitals participated with their regional stroke service in KNCN. Most coordinators of the regional stroke services have a background as specialised stroke nurse and are appointed at the hospital. These coordinators participate in the national learning network of KNCN. Important skills of coordinators include: networking, organising meetings with professionals and organisations, supporting the exchange and/or exchanging (new) knowledge, collecting data on quality of care, promoting and/or making agreements on collaboration, reflecting on current practice

and stimulating continuous quality improvement. Within the learning network, coordinators and KNCN identify and share ‘good practices’, guidelines and studies on specific topics in (integrated) stroke care. An online toolbox with this information and national meetings are available for all coordinators. An online forum facilitated communication between coordinators. In addition, educational courses on specific topics, such as ‘creating good collaboration within the stroke services’ are organised.

With the objective of encouraging continuous quality improvement, stroke services can participate in KNCNs self-assessment study to get insights into their own performance compared with earlier measuring moments and other stroke services. In the current study, we explored the results of KNCNs self-assessment studies in 2012, 2015 and 2019. The primary objective of our study was to establish which integrated care activities were implemented within the Dutch stroke services in 2012, 2015 and 2019. In addition, since a knowledge gap on the best size or scale for the governance of integrated stroke services exists [15], we explored whether implementation of integrated care activities was associated with characteristics of the stroke services involved in order to get more insight what organisational features might benefit or hamper the coordination and collaboration of integrated stroke care. A deeper insight in these relations are relevant; some studies indicate a relationship between the quality of integrated care networks and the volume of target group and/or the amount of collaborative partners [15].

METHODS

STUDY DESIGN

All stroke services networks participating in KNCN, 72 (2012), 59 (2015) and 55 (2019), were invited for the study. Although some stroke services (e)merged during the years, our study could be regarded as a cohort study. Coordinators of all stroke services networks were invited to fill out an online self-assessment questionnaire on behalf of the stroke services network. Coordinators could discuss the questions with stakeholders or complete the questionnaire by themselves. This was up to the coordinators. Participation was voluntary. A short instruction about the self-evaluation tool was provided. The invitations were sent in February/March of the respective years; a reminder was sent to coordinators who did not respond initially. Data collection ended in April. Ethical approval from a medical ethics committee was not needed under Dutch law as this was an observational study among care professionals only.

SELF-ASSESSMENT QUESTIONNAIRE

Coordinators were requested to provide general characteristics about their regional stroke services network and about implemented integrated care activities within the network. The Development Model for Integrated Care

(DMIC) [16, 17] served as a basis for the self-assessment questionnaire (in Dutch). The same questionnaire was used in all three years. After a brief explanation, the coordinators were asked to answer 97 questions about activities related to integrated stroke care, answering “yes”, “no” or “don’t know” (see Appendix 1). The DMIC is a validated model and consists of 97 activities [16, 17] grouped into nine clusters:

1. Interprofessional teamwork: professionals collaborating and working in well-organised multidisciplinary teams in the care network to provide care for a well-described client group (3 items);
2. Roles and tasks: clarity about the expertise, roles and tasks of professionals in the stroke service with effective collaboration at all levels, with new partners and by allocating coordinating roles (8 items);
3. Patient-centeredness: integrated care and information flows are tailored to the needs and characteristics of patients. This also includes self-management support (10 items);
4. Commitment: collaborative commitment and goals in the care chain with clearly defined goals and collaborative targets (12 items);
5. Transparent entrepreneurship: conditions for innovation and leadership responsibilities for performance achievement and joint financial agreements covering the integrated care (7 items);
6. Result-focused learning: a learning climate aiming for continuous improvement in the results of the stroke service (14 items);
7. Quality of care: the design of a multidisciplinary care pathway throughout the care service, based on guidelines and with patient representatives’ involvement (including clients’ needs assessment) (7 items);
8. Delivery system: logistics and coordination procedures for streamlining the care process within the stroke service (e.g. agreements, procedures and tools) (18 items);
9. Performance management: measurement and analyses of the results of the care as delivered (performance targets, quality indicators, analysis of mistakes and near misses, feedback, improvement activities) (18 items).

Furthermore, the DMIC distinguishes four phases of development: (1) the initiative and design phase; (2) the experimental and execution phase; (3) the expansion and monitoring phase; and (4) the consolidation and transformation phase. Each of development is described and characterised by the ten most phase-relevant activities. The development phase of a network is assessed by the overall description and the presence of activities measured by the DMIC. A phase is considered implemented if at least seven out of ten phase specific items are met. (see Appendix 2).

DATA ANALYSES

Descriptive analyses were conducted to study the characteristics of integrated stroke care services and the implementation of integrated stroke care activities. Stroke service cluster scores were calculated as the proportion (completed = 'yes') of implemented elements per cluster among all stroke services. The reported 'don't know' answers for implemented elements were regarded as 'not completed', as coordinators filled out the questionnaire in collaboration with stakeholders of the regional network. The associations between the cluster scores and characteristics of services were examined by means of linear regression analysis. Stepwise forward selection (probability-of-F-to-enter ≤ 0.05) was used in order to analyse the association of characteristics

and the clusters cautiously as we had no theoretical assumptions or model for the associations between characteristics and clusters. SPSS 25.0 was used for the data analyses.

RESULTS RESPONDENTS

Totals of 67 (2012), 53 (2015) and 47 (2019) stroke services responded to the invitation for the study. Over the years, the response rate remained high and varied from 93.1% (2012) to 85.5% (2019). Unfortunately, completed data about the general characteristics of stroke services was only partially available in 2012 due to a shortcoming in data collection ([Table 1](#)).

	2012	2015	2019
Number of stroke services invited	72	59	55
Number of stroke serviced responded (response rate in %)	67 (93.1)	53 (89.8)	47 (85.5)
Characteristics of regional stroke services networks			
Age, median (n)	9 (41) ¹	12 (53)	15 (46)
– Range	1–16	3–20	1–24
Number of organisations involved, median (n)	5 (40)	6 (53)	7 (47)
– Range	2–19	2–19	2–23
– 0–4 organisations, %	32.5	26.4	23.4
– 5–7 organisations, %	42.0	41.5	31.9
– 8–9 organisations, %	15.5	11.3	21.3
– 10 organisations or more, %	10.0	20.8	23.4
Volume of stroke patients, median (n)	364 (41)	450 (53)	499 (46)
– Min-Max volume	100–1552	79–1650	80–1200
– 1–300, %	34.1	20.8	21.7
– 301–599, %	43.9	56.6	41.3
– 600–899, %	19.6	18.9	15.3
– 900 or more, %	2.4	3.8	21.7
Coordinator of the regional stroke services network			
– Working hours per week, median (n)	8.0 (37)	8.0 (50)	8.0 (39)
– Range hours per week	1–24	1–24	3–24
– 1–4 hours per week, %	27.0	18.0	25.6
– 5–8 hours per week, %	27.1	45.3	38.5
– 9–12 hours per week, %	24.3	17.0	12.8
– 13 hours or more per week, %	21.6	15.1	23.1
– Appointed (% yes)	91.6	100.0	93.5
Working groups (% yes)	86.6	94.3	95.7
Formal agreements on cooperation (% yes)	80.6	86.8	82.4
Regular meetings with steering group (% yes)	77.4	90.6	73.9

Table 1 Characteristics of the study population in 2012, 2015 and 2019.

¹Because no data were collected in 2012, 26 stroke services did not receive questions on characteristics.

CHARACTERISTICS OF THE REGIONAL STROKE SERVICES NETWORKS

In 2012, most of the stroke services responding had existed for nine years. The median age of the regional stroke services networks increased proportionally over the years. By 2019, most networks had existed for 15 years. During these years, the number of organisations involved in the networks increased; in 2012, most regional stroke services networks consisted of five organisations and by 2019 this had increased to seven organisations (partly due to the merging of some networks). In addition, the median volume of stroke patients ('incidence') in the regional network increased: 364 patients in 2012 to 499 patients in 2019. Although over the years most coordinators were appointed for eight hours per week, the portion of coordinators appointed for more than eight hours per week was less in 2019 (35.1%) compared to 2012 (45.9%). In 2019, all stroke services networks had both a steering group and a working group. Three quarters of the stroke services networks had regular meetings with a steering group ([Table 1](#)).

IMPLEMENTING INTEGRATED CARE ACTIVITIES

The regional stroke services networks implemented most activities in two clusters 'Interprofessional teamwork' (3 items) and 'Roles and tasks' (8 items) ([Table 2](#)). In 2019, the networks had implemented 88.7% of the

activities for 'Interprofessional teamwork' and 85.6% of the activities for 'Roles and tasks'. Between 2012 and 2019, a statistically significant increase in the mean proportion of activities integrated was found for 'Transparent entrepreneurship', 'Roles and tasks', 'Patient-centeredness' and 'Commitment' ('Transparent entrepreneurship' (MD = 18.0% F(1) = 10.693, p = 0.001), 'Roles and tasks' (MD = 14.0% F(1) = 9.255, p = 0.003), 'Patient-centeredness' (MD = 12.9% F(1) = 9.255, p = 0.003), and 'Commitment' (MD = 11.2% F(1) = 4.982, p = 0.028). These improvements were mostly made during 2012 and 2015. After 2015, a reduction in implemented activities within the clusters occurred in the networks. This reduction was statistically not significant. 'Performance management' decreased by nine percentage points from the average score in 2015 (F(1) = 4.068, p = 0.046).

Most networks moved on to a later phase of the Development Model for Integrated Care (χ^2 (6) = 15.669, p = 0.016, n = 167). In 2019, 71 percent of the networks had reached one of the last two phases ('Expansion and monitoring phase' and 'Consolidation and transformation phase') ([Table 3](#)).

ASSOCIATION BETWEEN CHARACTERISTICS OF NETWORKS AND CLUSTER SCORES

The exploration of the association between characteristics of stroke services networks and cluster scores confirmed a positive correlation between the age of the regional

	2012	2015	2019	2012–2015	2015–2019	2012–2019
	mean % (95% CI)	mean % (95% CI)	mean % (95% CI)	mean difference F(df = 1)/p	mean difference F(df = 1)/p	mean difference F(df = 1)/p
	n = 67	n = 53	n = 47	n = 120	n = 100	n = 114
Interprofessional teamwork (3 items)	85.6 (79.7–91.5)	93.1 (88.9–97.3)	88.7 (82.8–94.5)	7.5 F = 3.924 p = 0.050	-4.4 F = 1.581 p = 0.212	3.1 F = 0.518 p = 0.473
Roles and tasks (8 items)	71.6 (65.6–77.7)	88.7 (84.4–92.9)	85.6 (78.8–92.4)	17.1 F = 19.097 p = 0.000	-3.1 F = 0.611 p = 0.436	14.0 F = 9.255 p = 0.003
Patient-centeredness (10 items)	59.0 (53.4–64.7)	71.0 (65.1–76.9)	71.9 (65.7–78.1)	12.0 F = 8.518 p = 0.004	-0.9 F = 0.045 p = 0.842	12.9 F = 9.246 p = 0.003
Commitment (12 items)	58.7 (52.5–65.0)	75.5 (69.5–81.4)	69.9 (61.9–77.8)	16.8 F = 14.515 p = 0.000	-5.6 F = 1.319 p = 0.254	11.2 F = 4.982 p = 0.028
Transparent entrepreneurship (7 items)	49.5 (42.4–56.6)	65.0 (59.0–71.0)	67.2 (59.1–75.2)	15.5 F = 10.375 p = 0.002	-2.2 F = 0.202 p = 0.654	18.0 F = 10.693 p = 0.001
Result-focused learning (14 items)	58.5 (52.2–64.8)	67.1 (61.6–72.6)	67.4 (61.0–73.8)	8.6 F = 3.938 p = 0.050	0.3 F = 0.005 p = 0.945	8.9 F = 3.675 p = 0.058
Quality of care (7 items)	53.5 (48.5–58.6)	66.3 (60.7–71.9)	61.4 (54.5–68.3)	12.8 F = 11.427 p = 0.001	4.9 F = 2.466 p = 0.120	7.9 F = 3.585 p = 0.061
Delivery system (18 items)	54.3 (49.7–59.0)	67.5 (62.1–72.8)	60.3 (53.4–67.2)	13.2 F = 13.962 p = 0.000	-7.2 F = 2.831 p = 0.096	6.0 F = 2.221 p = 0.139
Performance management (18 items)	52.2 (46.5–57.8)	67.1 (61.6–72.6)	58.0 (50.6–65.4)	14.9 F = 13.723 p = 0.000	-9.1 F = 4.068 p = 0.046	5.8 F = 1.596 p = 0.209

Table 2 Completed activities within the nine clusters of integrated care activities in 2012, 2015 and 2019¹.

¹Cluster scores were calculated as proportions of the activities implemented.

PHASE OF DEVELOPMENT	2012 (N = 67)	2015 (N = 53)	2019 (N = 47)
Initiative and design phase	37.3	34.0	8.5
Experimental and execution phase	9.0	7.5	21.3
Expansion and monitoring phase	35.8	35.8	48.9
Consolidation and transformation phase	17.9	22.6	21.3

Table 3 Stroke services per phase of development based on the Development Model for Integrated Care in 2012, 2015 and 2019^{1,2} (in %).

¹The phase of development is assessed by having implemented at least seven of the top-ten relevant items for a certain phase of development (see Appendix 2).

²Pearson's χ^2 (6) = 15.669, $p = 0.016$, $n = 167$.

stroke services networks and some clusters ('Roles and tasks' (B = 1.387 CI = 0.763–2.011 $p = 0.000$), 'Commitment' (B = 1.625 CI = 0.862–2.388 $p = 0.000$), 'Transparent entrepreneurship' (B = 1.912 CI = 1.152–2.677 $p = 0.000$), 'Result-focused learning' (B = 1.491 CI = 0.784–2.199 $p = 0.004$), 'Delivery system' (B = 1.402 CI = 0.754–2.050 $p = 0.000$) and 'Performance management' (B = 1.495 CI = 0.31–2.159 $p = 0.000$). Besides, the number of organisations involved is also associated with further implementation on 'Transparent entrepreneurship' (B = 1.607 CI = 0.534–2.680 $p = 0.004$) and 'Quality of care' (B = 1.088 CI 0.071–2.106 $p = 0.036$). A small but negative association was found between the volume of patients and performing the activities in two clusters, 'Interprofessional teamwork' (B = -0.015, CI = -0.028 to -0.003, $p = 0.019$) and 'Patient-centredness' (B = -0.014, CI = -0.027 to -0.001, $p = 0.040$) (**Table 4**).

DISCUSSION

A unique national self-assessment cohort study among stroke services was conducted to establish which integrated activities were performed by Dutch regional stroke networks in 2012, 2015 and 2019. This first national longitudinal study revealed that stroke services had developed further during this period of time, stroke services implemented statistical significantly more integrated care activities in four out of nine clusters (i.e. transparent entrepreneurship, roles and tasks, patient-centredness, commitment). Networks with more years of experience in collaboration and networks with more stakeholders involved, turned out to be further developed in terms of maturity of integrated care. Simultaneously, a higher volume of patients seems to be associated with performing less activities on patient-centredness and interprofessional teamwork.

The results resemble other study findings in studies of integrated care networks for other patient groups [18, 19]. Notwithstanding these positive results, integrated stroke care services networks should pay attention to implementing activities that are more linked to continuous quality improvement of care processes

and outcomes of care (the 'Result-focused learning', 'Quality of care', 'Delivery system' and 'Performance management' clusters). The way activities in integrated care networks should be implemented needs to be tailored. Realistic evaluations paying attention to what works for whom in what circumstances and how may be helpful in this respect [20, 21]. Some improvement activities may relate to minor changes in practice, whereas others may reveal larger cultural or organisational issues that need to be addressed and hence have longer timespans [22]. In addition to measuring the level of implementation of integrated activities it is necessary to also measure outcomes of integrated care to understand how outcomes can be improved. This is challenging since, among other things, data collection for further development of integrated care networks is hindered by the use of multiple registries in the organisations involved and the absence of a well-structured approach to systematically collect relevant and meaningful outcome data. A comprehensive quality improvement programme with a focus on continuous evaluation of the context, mechanisms and outcomes of integrated stroke care would be regarded helpful in enabling the provision of patient-centred, integrated care services [23, 24].

LEADERSHIP BY INTEGRATED CARE NETWORK COORDINATORS

A review on leadership in integrated care networks by Mitterlechner [25] underlines the importance of care coordinators in moving a network forward besides other factors such as network governance, trust and organisational structures. Integrated care network coordinators are needed for promoting collaboration in the network by encouraging communication, by creating inter-organisational linkages, by gathering stakeholders for problem solving and by facilitating the involvement of relevant parties. Within the regional networks, coordinators and stakeholders are then still able to evaluate the collaboration, processes and outcomes of integrated stroke care in their networks in a

	YEAR	R ²	AGE OF REGIONAL STROKE SERVICES NETWORK	ORGANISATIONS INVOLVED	VOLUME OF STROKE PATIENTS, LAST YEAR
Inter-professional teamwork (3 items)	All	0.045			B = -0.015 -0.028 to -0.003 <u>p = 0.019</u>
Roles and tasks (8 items)	2012	0.135	B = 2.074 0.247 to 3.902 <u>p = 0.027</u>		
	2019	0.140	B = 0.896 0.144 to 0.165 <u>p = 0.021</u>		
	All	0.137	B = 1.387 0.763-2.011 <u>p = 0.000</u>		
Patient-centeredness (10 items)	2019	0.118			B = -0.021 -0.041 to -0.002 <u>p = 0.035</u>
	All	0.092	B = 1.005 0.320 to 1.691 <u>p = 0.004</u>		B = -0.014 -0.027 to -0.001 <u>p = 0.040</u>
Commitment (12 items)	2015	0.098			B = 1.709 0.202 to 3.217 <u>p = 0.27</u>
	2019	0.227	B = 1.729 0.652 to 2.806 <u>p = 0.02</u>		
	All	0.160	B = 1.625 0.862 to 2.388 <u>p = 0.00</u>	B = 1.302 0.229 to 2.375 <u>p = 0.018</u>	
Transparent entrepreneurship (7 items)	2012	0.274	B = 2.776 0.656 to 4.897 <u>p = 0.012</u>	B = 3.454 0.605 to 6.304 <u>p = 0.019</u>	
	2015	0.085	B = 1.810 0.684 to 2.935 <u>p = 0.002</u>	B = 1.588 0.080 to 3.096 <u>p = 0.039</u>	
	2019	0.228	B = 1.914 1.152 to 2.677 <u>p = 0.00</u>	B = 1.607 0.534 to 2.680 <u>p = 0.004</u>	
	All	0.213	B = 1.914 1.152 to 2.677 <u>p = 0.00</u>	B = 1.607 0.534 to 2.680 <u>p = 0.004</u>	
Result-focused learning (14 items)	2019	0.173	B = 1.344 0.351 to 2.336 <u>p = 0.009</u>	B = 1.154 0.158 to 2.150 <u>p = 0.024</u>	
	All	0.155	B = 1.491 0.784 to 2.199 <u>p = 0.004</u>		
Quality of care (7 items)	2015	0.111		B = 1.705 0.305 to 3.105 <u>p = 0.018</u>	
	All	0.035		B = 1.088 0.071 to 2.106 <u>p = 0.036</u>	
Delivery system (18 items)	2019	0.204	B = 1.442 0.478 to 2.406 <u>p = 0.004</u>		
	All	0.131	B = 1.402 0.754 to 2.050 <u>p = 0.000</u>		
Performance management (18 items)	2019	0.294	B = 1.754 0.836 to 2.673 <u>p = 0.000</u>		
	All	0.140	B = 1.495 0.831 to 2.159 <u>p = 0.000</u>		

Table 4 Regression analysis with forward selection¹ on completed integrated activities within the nine clusters and characteristics of integrated stroke services within and over the years (B, CI, p).

¹Criterion: probability-of-F-to-enter \leq 0.050.

²Only the statistically significant correlations are presented. No correlations with ‘working hours of the coordinator’ were found.

practical and participatory way and support continuous reflective learning [19]. The KNCN learning network gives coordinators of regional stroke services networks the opportunity to ‘teach each other and to learn from each other’ which could also benefit the improvement of integrated care activities and learning between services.

SCALE OF THE REGIONAL STROKE SERVICES NETWORK

After 2015, an apparent reduction in completed activities within the clusters took place in the networks, due to some regional networks merging. A significant decrease in the

mean score for 'Performance management' was found. The association of integrated care activities with the volume of patients (small and negative) and the number of organisations involved (positive) seems interesting in this respect. Adding more organisations to the networks apparently encourages structured collaboration on integrated care in the region. On the other hand, a higher volume of patients could impede 'patient-centredness' and regional networks should consider which patients (or groups of patients) should be their target groups and what is the most suitable scale for stroke services. The relationship between scale and patient-centeredness remains mainly unexplored in integrated care [15]. Patient-centredness in integrated care could include activities concerning the design of processes of care, training of professionals, communication and information exchange and patient involvement in care [26].

Since 2018, the Dutch government has been promoting the transfer of care and the collaboration between stakeholders within localities by means of a national programme called 'The right care, at the right place, at the right time, with the right resources'. This movement favours regional collaboration, and therefore increases the need for effective implementation of integrated stroke care services within the regional networks at both the local and interregional levels especially when shortages of resources (e.g. staff, money) are experienced [15]. Further studies into the relationships of the organisation and scale of integrated care networks with patient-reported experiences and outcomes are needed.

STRENGTHS AND LIMITATIONS AND SUGGESTIONS FOR FURTHER RESEARCH

The strengths of the current study are the repeated measurements over a period of seven years by a validated model and the high response rate of stroke services. The DMIC is a model that helps to create an understanding of the presence of the activities as part of the delivery of integrated care services [27]. Although the DMIC was developed in a former decade, the model seems still valid, including the most relevant activities for an integrated care network. The model is still applied in recent studies and considered relevant in a diversity of countries and contexts, including Parkinson Networks in Germany, integrated services networks in Canada and diabetes networks in the Netherlands [19, 28–29]. However, this study also has some limitations. First, as this study uses a self-assessment method, the answers given are subjective and open to bias. Coordinators may have had different frames of reference or interpretations of the integrated activities or the level to which they have been implemented. Second, the questionnaires were completed by the coordinator on behalf of the stroke

services networks and no data is available about whether and how coordinators involved stakeholders of the regional networks in providing their answers. Thirdly, the DMIC does not involve an assessment of the outcomes of the integrated care services. This is a recommendation for further research. Fourth, the relationships between the characteristics (e.g. scale), development of services and outcomes need to be further addressed in future research. In our study, a regression analysis was used to explore the association between integrated care activities and characteristics of integrated stroke services. A better understanding is needed on the impact of features on performance. Finally, the study results are primarily valid for the Dutch healthcare system. Nonetheless, the Action Plan for Stroke in Europe 2018–2030 formulates targets for integrated stroke care that are in line with the results of our study: 1) someone should be responsible for stroke quality improvement in the region; 2) systems for assessing/accrediting stroke services are needed with peer support for quality improvement; and 3) patient-reported and clinical outcomes covering both hospital and community care should be collected to improve healthcare [30].

CONCLUSION

An increase was found in the development of integrated care activities within the regional stroke networks in the Netherlands, from 2012 until 2019. Further, experience in collaboration in the networks benefits the performance of integrated care activities. In the context of the Netherlands, stroke services with longer collaboration timespans do result in improved development phases, as defined by the DMIC, and a more comprehensive range of elements of integrated care. However, after two decades of implementation of stroke services, there is room for further improvement. This illustrates the long-term commitment that is needed if these complex programmes or strategies are to provide added value.

DATA ACCESSIBILITY STATEMENT

All data are archived by the first author (HVP).

ADDITIONAL FILES

The additional files for this article can be found as follows:

- **Appendix 1.** Overview Over Integrated Activities Asked for. DOI: <https://doi.org/10.5334/ijic.5611.s1>
- **Appendix 2.** Development Model for Integrated Care – Description of development phases. DOI: <https://doi.org/10.5334/ijic.5611.s2>

ETHICS AND CONSENT

No ethical approval from a medical ethics committee was needed under Dutch law as this was an observational study among care professionals only.

ACKNOWLEDGEMENTS

The authors would like to acknowledge all board members of Dutch Knowledge Network of Stroke Services (KNCN). We express special thanks to former chair and grounding father George Beusmans MD PhD, who passed away in 2020.

REVIEWERS

Prof Zhanchun Feng, School of Medicine and Health Management, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

Ronald Uittenbroek PhD., Head of school, school of nursing, Health and social care dept., Windesheim university of applied sciences, The Netherlands.

FUNDING INFORMATION

The project was financed by contributions of the regional networks to membership of the Dutch Knowledge Network of Stroke Services (KNCN). The regional networks had no role in the analysis of interpretation of data or in writing of the manuscript. Data analysis was financed by EnCorps.


COMPETING INTERESTS


The authors have no competing interests to declare.

AUTHOR CONTRIBUTIONS

HVP, MB, LvD, MK, HK initiated the evaluation study in 2019. HVP, NZ, MB, EvW were involved in the data collection in 2019. MM is the architect of the theoretical model and the questionnaire. HVP analysed the data. NZ, HK, HJMV, MM checked all data analysis. All authors read and approved the final manuscript.

AUTHOR AFFILIATIONS

Helene R. Voogdt-Pruis  orcid.org/0000-0001-7801-3897
University Medical Centre Utrecht, Julius Global Health, Utrecht University, PO Box 85500, 3508 GA Utrecht, Netherlands;
Encorps, 1214 ND, Hilversum Netherlands

Nick Zonneveld  orcid.org/0000-0003-0902-1927
Tilburg University, TIAS School for Business and Society, PO Box

90153, 5000 LE Tilburg, Netherlands; Vilans, National Centre of Expertise in Long-term Care, PO Box 8228, 3503 RE Utrecht, Netherlands

Monique Bergsma

Stroke Knowledge Network Netherlands, Oudlaan 4, 3515 GA Utrecht, Netherlands

Elize van Wijk

Vilans, National Centre of Expertise in Long-term Care, PO Box 8228, 3503 RE Utrecht, Netherlands

Henk Kerkhoff

Albert Schweitzer Hospital, Department of Neurology, PO Box 444, 3300 AK Dordrecht, Netherlands

Luikje van der Dussen

Stroke Knowledge Network Netherlands, Oudlaan 4, 3515 GA Utrecht, Netherlands

Maartje Kuijpers

Stroke Knowledge Network Netherlands, Oudlaan 4, 3515 GA Utrecht, Netherlands

Hubertus J. M. Vrijhoef

 orcid.org/0000-0002-5539-4671

Stroke Knowledge Network Netherlands, Oudlaan 4, 3515 GA Utrecht, Netherlands

Mirella M. N. Minkman

 orcid.org/0000-0003-4922-5983

Tilburg University, TIAS School for Business and Society, PO Box 90153, 5000 LE Tilburg, Netherlands; Vilans, National Centre of Expertise in Long-term Care, PO Box 8228, 3503 RE Utrecht, Netherlands

REFERENCES

1. **GBD (2016) Stroke Collaborators.** Global, regional, and national burden of stroke, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol*, 2019; 48: 439–458.
2. **van Eeden M, van Heugten C, van Mastrigt GAPG.** The burden of stroke in the Netherlands: estimating quality of life and costs for 1 year poststroke. *BMJ Open*, 2015; 5: e008220. DOI: <https://doi.org/10.1136/bmjopen-2015-008220>
3. **KNCN.** Healthcare standard integrated stroke. Utrecht: Dutch Knowledge Network of Stroke Services; 2012. <https://www.kennisnetwerkcva.nl/wp-content/uploads/2018/09/Zorgstandaard-CVA-TIA-printversie.pdf> (accessed 5 May 2020).
4. **WHO.** Integrated care models: an overview; 2016. <http://www.euro.who.int/>.
5. **Auscha C.** Barriers to the Integration of Care in Inter-Organisational Settings: A Literature Review. *Int J Integr Care*, 2018; 18(1): 5. Published 2018 Jan 16. DOI: <https://doi.org/10.5334/ijic.3068>
6. **Harteraad.** Kwaliteitscriteria CVA zorg vanuit patiëntperspectief; 2018.
7. **Voogdt-Pruis HR, Ras T, van der Dussen L, et al.** Improvement of shared decision making in integrated stroke care: a before and after evaluation using a questionnaire survey. *BMC Health Serv Res*, 2019; 19(1): 936. Published 2019 Dec 5. DOI: <https://doi.org/10.1186/s12913-019-4761-2>
8. **van der Linden BA, Spreeuwenberg C, Schrijvers AJ.** Integration of care in The Netherlands: the development

- of transmural care since 1994. *Health Policy*, 2001 Feb; 55(2): 111–20. DOI: [https://doi.org/10.1016/S0168-8510\(00\)00125-1](https://doi.org/10.1016/S0168-8510(00)00125-1)
9. **Rosendal, H, Wolters, CAM, Beusmans G**, et al. Stroke service in The Netherlands: an exploratory study on effectiveness, patient satisfaction and utilisation of healthcare. *International journal of integrated care*, 2002; 2:e17. DOI: <https://doi.org/10.5334/ijic.50>
 10. **IHI**. The breakthrough series: IHI's collaborative model for achieving breakthrough improvement. In *IHI Innovation Series white paper*; 1995. Boston: Institute for Healthcare Improvement. www.ihio.org.
 11. **Kilo CM**. A framework for collaborative improvement: Lessons from the Institute for Healthcare improvement's Breakthrough series. *Quality Management in Health Care*, 1998; 6. DOI: <https://doi.org/10.1097/00019514-199806040-00001>
 12. **Minkman MMN, Schouten LMT, Huijsman R, Splunteren van PT**. Integrated care for patients with a stroke in the Netherlands; results and experiences from a Breakthrough Collaborative Improvement project. *International Journal of Integrated Care*, March 2005; 5(23). ISSN 1568-4156.
 13. **Huijsman R, Rasch P**. Borgen, sturen en innoveren. In: *De beste zorg bij CVA. Handreiking voor het verbeteren van de CVA zorgketen*. CBO; 2005.
 14. **Miller R, Stein KV**. The Odyssey of Integration: Is Management its Achilles' Heel? *Int J Integr Care*, 2020; 20(1): 7. Published 2020 Feb 26. DOI: <https://doi.org/10.5334/ijic.5440>
 15. **Minkman MMN**. Suitable Scales; Rethinking Scale for Innovative Integrated Care Governance. *International Journal of Integrated Care*, 2020; 20(1): 1. DOI: <https://doi.org/10.5334/ijic.5468>
 16. **Minkman M**. Developing integrated care. Towards a development model for integrated care. Thesis. University of Rotterdam; 2012. ISBN-978-90-13-10026-6801-300.
 17. **Minkman MMN, Ahaus K, Fabbriotti I**, et al. Quality management model for integrated care: results of a Delphi and Concept Mapping study. *International Journal on Quality in Health Care*, 2009 Feb; 21(1): 66–75. DOI: <https://doi.org/10.1093/intqhc/mzn048>
 18. **Vat LE, Middelkoop I, Buijck BI**, et al. The Development of Integrated Stroke Care in the Netherlands a Benchmark Study. *International Journal of Integrated Care*, 2016; 16(4): 12. DOI: <https://doi.org/10.5334/ijic.2444>
 19. **Zonneveld N, Vat LE, Vlek H**, et al. The development of integrated diabetes care in the Netherlands: a multiplayer self-assessment analysis. *BMC Health Services Research*, 2017; 17: 219. DOI: <https://doi.org/10.1186/s12913-017-2167-6>
 20. **Busetto L, Luijckx KG, Elissen AMJ**, et al. Context, mechanisms and outcomes of integrated care for diabetes mellitus type 2: a systematic review. *BMC Health Serv Res*, 2016; 16: 18. DOI: <https://doi.org/10.1186/s12913-015-1231-3>
 21. **Goodwin N**. Improving Integrated Care: Can Implementation Science Unlock the 'Black Box' of Complexities? *International Journal of Integrated Care*, 2019; 19(3): 12, 1–3. DOI: <https://doi.org/10.5334/ijic.4724>
 22. **SUSTAIN**. Sustainable Tailored Integrated Care for Older People in Europe: The Roadmap; 2019. Available at: <https://www.sustain-eu.org/wp-content/uploads/sites/4/2019/03/SUSTAIN-Roadmap.pdf>.
 23. **Zipfel N, van der Nat PB, Rensing BJWM**, et al. The implementation of change model adds value to value-based healthcare: a qualitative study. *BMC Health Services Research*, 2019; 19: 643. DOI: <https://doi.org/10.1186/s12913-019-4498-y>
 24. **Wankah P, Guillette M, Lemaitre T**, et al. Challenges in measuring integrated care models: International knowledge and the case of Quebec. *International Journal of Care Coordination*; 2020. DOI: <https://doi.org/10.1177/2053434520945087>
 25. **Mittlechener M**. Leadership in integrated care networks; a literature review and opportunities for future research. *International Journal of Integrated Care*, 2020; 20(3): 6, 1–14. DOI: <https://doi.org/10.5334/ijic.5420>
 26. **Stoop A, Lette M, Ambugo EA**, et al. Improving person-centredness in integrated care for older people: Experiences from thirteen integrated care sites in Europe. *International Journal of Integrated Care*, 2020; 20(2): 16, 1–16. DOI: <https://doi.org/10.5334/ijic.5427>
 27. **Bautista MA, Nurjono M, Lim YW**, et al. Instruments Measuring Integrated Care: A Systematic Review of Measurement Properties. *Milbank Q*, 2016; 94(4): 862–917. DOI: <https://doi.org/10.1111/1468-0009.12233>
 28. **Longpré C, Dubois C-A**. Implementation of integrated services networks in Quebec and nursing practice transformation: convergence or divergence? *BMC health services research*, 2015; 15(1): 84. DOI: <https://doi.org/10.1186/s12913-015-0720-8>
 29. **Loewenbrück KF, Stein DB, Amelung VE**, et al. Parkinson Network Eastern Saxony (PANOS): Reaching Consensus for a Regional Intersectoral Integrated Care Concept for Patients with Parkinson's Disease in the Region of Eastern Saxony, Germany. *Journal of Clinical Medicine* [Internet], 2020 Sep [cited 2020 Nov 9]; 9(9): 2906. Available from: <https://www.mdpi.com/2077-0383/9/9/2906>. DOI: <https://doi.org/10.3390/jcm9092906>
 30. **Norrving B, Barrick J, Davalos A**, et al. Action Plan for Stroke in Europe 2018–2030. *European stroke journal*, 2018; 3(4): 309–336. DOI: <https://doi.org/10.1177/2396987318808719>

TO CITE THIS ARTICLE:

Voogdt-Pruis HR, Zonneveld N, Bergsma M, van Wijk E, Kerkhoff H, van der Dussen L, Kuijpers M, Vrijhoef HJM, Minkman MMN. Optimising Integrated Stroke Care in Regional Networks: A Nationwide Self-Assessment Study in 2012, 2015 and 2019. *International Journal of Integrated Care*, 2021; 21(3): 12, 1–11. DOI: <https://doi.org/10.5334/ijic.5611>

Submitted: 10 September 2020 Accepted: 08 September 2021 Published: 20 September 2021

COPYRIGHT:

© 2021 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

International Journal of Integrated Care is a peer-reviewed open access journal published by Ubiquity Press.

