

---

## POSTER ABSTRACT

### A quasi experimental before and after study of a transitional care programme for older adults in the area of the IJsselland Hospital

18<sup>th</sup> International Conference on Integrated Care, Utrecht, 23-25 May 2018

Wilma van der Vlegel-Brouwer<sup>1</sup>, Paul Breman<sup>2</sup>, Lois Thomas<sup>3</sup>, Lyndsey Macphail<sup>3</sup>

1: IJsselland hospital, The Netherlands;

2: University of Applied Sciences Utrecht, The Netherlands;

3: Uclan University of Central Lancashire, United Kingdom

---

**Background:** In the Netherlands increasing numbers of older adults are living at home. Older people being discharged from hospital, at high risk of functional decline, need support to enable them to continue to live at home and prevent re-admission or admission to residential care. A transitional care program developed in Amsterdam, "the Transitional Care Bridge", is embraced by care insurers. The programme has been adopted by the IJsselland hospital together with six home care organizations. In this programme the identification of seniors at risk starts in the Emergency Department. Patients at high risk fill out a minimal Comprehensive Geriatric Assessment CGA form. The geriatric team visits the frailest patients. If necessary, patients are followed up for six months by the Community nurse, with the first visit the day before discharge from hospital.

**Methods:** Two groups of 50 older patients are compared over a three month period in a 'before and after' quasi-experimental study. Patients were asked to fill out a questionnaire based on the TOPICS-MDS, containing several validated instruments, before discharge from hospital, and after one and three months post-discharge. Hypotheses are the programme will have an effect on mortality, length of stay, self-management, self-rated health and quality of life, and use of formal and informal care. These hypotheses were tested and outcomes of the oldest old >85 years compared with the average old 75-85 years. Self-management abilities were assessed by the patient and the CN in the intervention group by filling out the Self Sufficiency Monitor. This instrument based on the ten standardized items was validated in this study with Cronbach's Alpha 0,802.

**Results:** Although mortality and the length of stay dropped in the intervention group, no statistically significant difference was found. Chi square = 0,444, p = .505, T = 0.825, p = .411. No statistically significant difference was found in use of care after one and three months after discharge: GP visits, T = -1,252, p = .214, T = -0,767, p = .445; ED visits, T = -0,581, p = .562, T = -0,045, p = .964; readmissions, T = -0,560, p = .577, T = 0,119, p = .095; admission to care home, T = 0,000, p = 1.000, T = -0,891, p = .375; help from Community Nurse CN, Chi2 = 0,185, p = .667, Chi2 = 1,013, p = .314. No effect was found on informal care, as well as on risk of delirium, functional decline, self-rated health and quality of life. On self-

van der Vlegel-Brouwer; A quasi experimental before and after study of a transitional care programme for older adults in the area of the IJsselland Hospital

management no statistically significant differences were found,  $F = 1,933$ ,  $p = .150$ . The intervention group was however, statistically significant more frail after one and three months after discharge,  $T = -2.514$ ,  $p = .014$  and  $T = -3.911$ ,  $p = .000$ .

**Discussion:** Results are influenced by the small sample and the fact that the intervention sample was more frail. Which was probably due to the extra triage in the intervention group by the minimal CGA.

**Further research:** A process evaluation using qualitative approaches will be conducted to further explore delivery of the programme, and patients' and health professionals' perspectives on its usefulness.

---

**Keywords:** older adults; transitional care; mixed methods evaluation; before and after study; oldest old

---